



# Antioxidant Potential of Seven Myrtaceous Fruits

Kurt A Reynertson, Margaret J. Basile,  
and Edward J. Kennelly

## Abstract

Many fruits of the Myrtaceae have a rich history of use both as edibles and as traditional medicines in divergent ethnobotanical practices throughout the tropical and subtropical world. From South America to Southeast Asia, these fruits have been used for a wide variety of ailments, including cough, diabetes, dysentery, inflammation and ringworm. These same fruits are also used to make many food products. Based on information regarding ethnomedicinal use, known phytochemistry, fruit color, popularity as edibles and availability, the fruits of several edible species from the subtribe Eugeniinae have been selected for phytochemical analysis in an attempt to discover new antioxidants. The fruits of six species in this group have shown a strong antioxidant activity in the 1,1-diphenyl-2-picrylhydrazyl chemical assay. The UV absorbance spectrum of the most active compound in *Eugenia uniflora* L. indicates that it is a flavonoid. Polyphenolic compounds like flavonoids have an enormous range of biological activity and are known to inhibit oxidative damage in vivo better than the classical vitamin antioxidants. In plants, they protect against lipid peroxidation and UV damage that can affect tropical fruits growing under severe conditions including high heat and intense sunlight.

## Introduction

In an initial screening of over thirty edible tropical fruits, extracts of *Eugenia uniflora* L. (Myrtaceae) proved to be highly active in the 1,1-diphenyl-2-picrylhydrazyl (DPPH) antioxidant assay, and polyphenolic compounds are implicated in this activity. A literature search of the plants most closely allied to *E. uniflora* was therefore conducted based on the hypothesis that allied species may contain similar antioxidant compounds. The ethnomedicinal and ethnobotanical uses of these plants was correlated with known phytochemical data to ascertain a list of promising species for further research to find new potent antioxi-

dant fruits and compounds. Increasing understanding of the role oxidative stress plays in disease has made the search for antioxidants more important.

## Antioxidants and Disease

Oxidative damage in the human body plays an important causative role in disease initiation and progression (Yamaguchi *et al.* 1998). Oxidation of low-density lipoprotein (LDL) is a major factor in the promotion of heart disease and atherosclerosis (Frankel *et al.* 1993a, Jacob & Burri 1996, Steinberg 1997). Damaging free radicals and reactive oxygen species (ROS) are produced naturally through oxidative metabolism and have also been linked to some cancers (Frankel *et al.* 1993a, Jacob & Burri 1996). Damage is generally reduced by endogenous antioxidants, but additional protection is necessary, and nutritive elements from food are critical in disease prevention. Repeated damage caused by ROS throughout the span of a human life increases with time, and is a major cause of age-related cancers and other oxidatively-induced diseases.

## Correspondence

Kurt A. Reynertson and Edward J. Kennelly, The Graduate Center, City University of New York and Lehman College, Department of Biology, 250 Bedford Park Boulevard West, Bronx, NY 10468. U.S.A. allerslev@gmail.com, Edward.Kennelly@lehman.cuny.edu  
Margaret J. Basile, University of Miami School of Medicine, Department of Neurology, 1501 NW 9th Avenue, Miami, FL 33136. U.S.A.

**Ethnobotany Research & Applications 3:025-035 (2005)**

Diets high in fruits and vegetables and low in cholesterol and fats are inversely correlated with the incidence of coronary heart disease (CHD) and cancer (Hertog *et al.* 1993, Hertog *et al.* 1995, Knekt *et al.* 1996, Ness & Powles 1999, Pietta 2000, van Poppel *et al.* 1994). Natural antioxidants from fruits and vegetables provide a measure of protection that slows the process of oxidative damage, and are implicated as the protective constituents of these foods (Hertog *et al.* 1993, Hollman & Katan 1999, van Poppel *et al.* 1994). Research in natural antioxidants is becoming increasingly important both in understanding the beneficial aspects of plant foods and in improving the quality of fatty foods. Antioxidants are routinely used by the industry to prevent the oxidation of food in storage and inhibit rancidity. The well-known vitamin antioxidants in food include ascorbic acid, b-carotene and a-tocopherol. Many clinical and epidemiological studies have sought to demonstrate the efficacy of these vitamins in preventing a wide variety of diseases (Blumenthal *et al.* 2000, Giugliano 2000, Haeghele *et al.* 2000, Jacob & Burri 1996, Pellegrini *et al.* 2000, Scheen 2000). However, some of these studies failed to show significant antioxidative protection in vitro (Pellegrini *et al.* 2000, Scheen 2000), which suggests that vitamins obtained via whole food or by a balanced diet may be more effective than supplements, possibly through synergistic interactions with other compounds.


### Polyphenolic Antioxidants

Recent studies have shown that many flavonoids and related polyphenols are actually better antioxidants than vitamins (Frankel *et al.* 1993b, Pietta 2000, Vinson *et al.* 1999a). Fruits and vegetables are high in flavonoid content; flavonoids impart color and taste to flowers and fruits, and it is estimated that humans consume between a few hundred milligrams and one gram of flavonoids every day (Hollman & Katan 1999, Pietta 2000). Flavonoids appear in blood plasma at pharmacologically active levels after eating flavonoid-rich foods but do not accumulate in the body (Cao *et al.* 1998, Hollman & Katan 1999). Consuming flavonoids regularly increases longevity by reducing inflammation and contributing to the amelioration of atherosclerosis from CHD (Frankel *et al.* 1993a). The range of flavonoid biological activity is large; in addition to scavenging free radicals and ROS, flavonoid actions include anti-inflammatory, antiallergenic, antiviral, antibacterial, antifungal, antitumor, and antihemorrhagic (Formica & Regelson 1995, Slowing *et al.* 1996). Flavonoids also inhibit a number of enzymes, including aldose reductase, a-glucosidase, xanthine oxidase, monooxygenase, lipoxigenase and cyclooxygenase (Middleton & Kandaswami 1993, Yoshikawa *et al.* 1998). Plant polyphenols interact with LDL, enriching and protecting it from oxidation when entering the bloodstream. The so-called "French Paradox" refers to the fact that despite the high fat content of the French diet, there is a lower incidence of CHD in France than in countries where fat intake is similar. This has been

attributed to the high polyphenolic content of red wine and other fruits and vegetables (Burns *et al.* 2000, Frankel *et al.* 1993a).

There are over 4000 naturally occurring flavonoids in several subclasses. All have the same basic C6-C3-C6 phenolic carbon skeleton (Figure 1). Flavonoids are ubiquitous in the higher plants and play an ecological as well as physiological role. The anthocyanins are the most important flower and fruit pigments; they attract pollinators and seed dispersers and protect plant tissues from ultraviolet (UV) radiation damage. Some flavonoids act as antifeedants to herbivorous pests. The isoflavones are responsible for the chemical signaling involved in legumous root nodule formation (Dewick 1997, Harborne & Baxter 1999, Robinson 1991).

**Figure 1.** Basic structures of **A** a flavonoid and **B** an anthocyanidin.



Studies suggest that the antioxidant potential of phenolics is mainly due to their ability to act as reducing agents (Burns *et al.* 2000, Kähkönen *et al.* 1999, Vinson *et al.* 1999b). It is well established that the efficacy of flavonoids as antioxidants stems from the number and position of the hydroxyl substitutions on the basic structure; an increase in number of hydroxyl groups is directly correlated with increasing activity, and the 3',4' -dihydroxy substitution is significant (Cao *et al.* 1997, Rice-Evans *et al.* 1996, Rice-Evans *et al.* 1997).

Anthocyanins (Figure 1B) are the glycosides of anthocyanidins, and contribute greatly to the antioxidant properties of certain fruits. As pigments, they produce the orange, red and blue colors in fruits and flowers. The antioxidant anthocyanidin that colors blueberries and grapes bluish-red is delphinidin. Other known antioxidant anthocyanins include cyanidin (orange-red), pelargonidin (orange), malvidin (bluish-red) and peonidin (red) (Wang *et al.* 1997). Fruit color is therefore an important indicator of possible polyphenolic compounds.

### Chemotaxonomic Approach to Antioxidant Discovery

As early as 1897, Baker and Smith investigated the essential oils of Eucalyptus (Myrtaceae) and found a close connection between the chemistry of the oils and the taxonomy of the plants (Gibbs 1993). It has since been established that chemistry is very useful in plant systematics. Likewise, systematics can be used in the search for bioactive compounds, and flavonoids are considered excellent taxonomic guides (Bate-Smith 1963). While flavonoids are ubiquitous in the higher plants, certain subclasses of flavonoids can be taxa-specific. It is unusual for rare flavonoids to occur outside a group in which they have been discovered. Glycosidic combinations are proving to be highly specific within families, and some morphological characteristics can be linked to particular flavonoid patterns (Harborne 1963). Some methylation patterns occur only in certain families (Pierpoint 1986). Anthocyanins can occur in any part of a plant, and different parts of the same plant often have different anthocyanin pigments. The glycosylation pattern is often consistent in a plant, and correlates with systematic information (Harborne 1963). Harborne found that species which do not conform to general glycosidic patterns are exceptional in other respects as well. Leaves and fruits tend to have simpler pigments than flowers, and there is an evolutionary trend toward more complex anthocyanin structures (Harborne 1963). Complex pigments with several glycosylations are more stable to light degradation and enzymatic attack. The evolutionary trend towards complexity is paralleled by a trend toward the blue color, which in flowers is related to the color preference of insect pollinators.

Anthocyanin content of fruits tends to increase as the fruit matures, becoming complexed with metals and other flavonoids (Pierpoint 1986, Silva 1997). The most common anthocyanidin is cyanidin, which occurs in 80% of permanently pigmented leaves, 69% of fruits and 50% of flowers. The next two most common anthocyanidins are delphinidin and pelargonidin. Delphinidin is wholly absent from some families, and abundant in others. As of 1995, only about 1000 species had been examined for anthocyanins, and only about one-fifth of those species have had the sugar groups fully described. This represents a tiny fraction of the 250,000 known species of angiosperm. Harborne believes that there must be a considerable number of new anthocyanin structures yet to be discovered (Harborne 1963, Pierpoint 1986).

The chemotaxonomic approach to antioxidant discovery includes reviewing species that have been phytochemically examined. Information can be extrapolated to allied plants. Species closely related to plants containing known polyphenolic antioxidants are likely to have similar polyphenolic constituents. Therefore, the phytochemistry of one plant may be used as a clue for a related plant. The hypothesis is that antioxidant activity may "run in the family."

### Plant Selection and Collection

Ethnobotanists regularly structure questionnaires to probe indigenous knowledge for the medicinal uses of plants. Most indigenous knowledge does not include a list of plants that help scavenge free radicals. Researchers looking for new antioxidants can, however, use ethnomedicinal information as a guide, paying special attention to plants that are used for illnesses or conditions that are ameliorated by compounds also linked to antioxidant activity. Polyphenolics are diverse in their biological activities, so ethnomedicinal information that hints at polyphenolic content could point to possible antioxidants.

For this study, the plants closely allied to *E. uniflora* were reviewed for ethnobotanical and phytochemical data. Several databases were queried, including NAPRALERT, Chemical Abstracts and Biological Abstracts. The number of NAPRALERT hits for each species studied is given in Table 1. Some of the species queried had little or no phytochemical or ethnobotanical data. Those that are widely used as food or medicine have often been phytochemically examined.

Several species were selected, and seven have been collected and tested: *Eugenia aggregata* (Velloso) Kiaersk., *E. foetida* Pers., *E. stipitata* McVaugh, *E. uniflora*, *Myrciaria cauliflora* (Mart.) O. Berg, *Syzygium jambos* (L.) Alston and *S. samarangense* (Blume) Merr. & L.M. Perry). Several other species have also been marked for testing, and analysis will be undertaken when fruits ripen. All are closely related species of the Myrtaceae subtribe Eugeniinae, and most are cultivated for their edible fruits. Some of these species are also used medicinally, although thorough phytochemical studies have not been done for each. Table 1 summarizes the known phytochemistry and ethnobotany of the fruits.

Several institutions in southern Florida dedicated to the propagation of tropical fruits generously permitted us to collect fruit. Collecting within the United States eliminates the need for international collection permits, and these institutions represent collections of plants that have been pre-selected and imported as edibles, medicinals or both. Fruit was collected from The Kampong (The National Tropical Botanical Gardens), The Broward County Rare Fruit and Vegetable Council Experimental Plot, the Fruit and Spice Park and The University of Florida Tropical Resource and Education Center. Fruits were shipped frozen to the laboratory, where they were kept at  $-20^{\circ}\text{C}$  until extraction. Voucher specimens have been deposited in the New York Botanical Garden herbarium.

### The Myrtle Family

The Myrtaceae is a well defined family, with leathery glandular leaves containing viscous aromatic terpenoid and

**Table 1.** Ethnobotanical and phytochemical information of Eugenioid fruits. Total number of NAPRALERT hits is given, along with antioxidant activity expressed as a DPPH IC50. A lower IC50 value indicates greater activity. Known vitamin antioxidants ascorbic acid and  $\alpha$ -tocopherol have IC50 values of 18.3mg/mL and 53.3mg/mL, respectively.


Species	Vernacular Name	Fruit Color	Ethnobotanical Information	Previous Phytochemical Work	Total	DPPH IC50 (mg/mL)
<i>Eugenia aggregata</i>	Rio Grande cherry	Reddish-purple	Brazil: eaten fresh, used for jams and jellies <sup>1</sup>	Catechin and epicatechin <sup>2</sup>	0	74.1
<i>Eugenia foetida</i>	Topiary bush	Purple	Hedge, topiary		0	15.9*
<i>Eugenia stipitata</i>	Araca-boi	Yellow	Desserts <sup>7</sup>	Terpenes, primarily sesquiterpenes <sup>3</sup>	0	79.0
<i>Eugenia uniflora</i>	Surinam cherry	Red to purple to black	Brazil: eaten fresh, used for desserts, liqueurs, wines (syrops and wines used medicinally); <sup>4</sup> astringent, high blood pressure <sup>5</sup>	Ascorbic acid, b-carotene, and a few sesquiterpenes <sup>6</sup>	235	19.6
<i>Myrciaria cauliflora</i>	Jaboticaba	Dark red to purple-black	Brazil: eaten fresh, used for jam, tarts, strong wine and liqueur, as a treatment for hemoptysis, asthma, diarrhea dysentery and chronic inflammation of the tonsils. <sup>7</sup>	Tannins <sup>7</sup>	0	35
<i>Syzygium jambos</i>	Rose apple	Yellow with slight blush	India: tonic for the brain and for liver problems, as an astringent, and digestive, <sup>8</sup> distilled to make rosewater. <sup>7</sup>	Terpenoids	282	247
<i>Syzygium samarangense</i>	Wax jambu	Pink to red	India: eaten fresh; <sup>9</sup> Malaya: greenish fruits are eaten raw with salt or cooked as a sauce <sup>7</sup>	Two flavonol glycosides as well as epigallocatechin 3-O-gallate, epicatechin 3-O-gallate, and samarangenin A and B. <sup>10</sup>	61	76.8

<sup>1</sup>Facciola 1998, <sup>2</sup>Reynertson unpublished, <sup>3</sup>Franco 2000, <sup>4</sup>Popenoe 1920, <sup>5</sup>Bandoni *et al.* 1972, <sup>6</sup>Duke 2000, Rucker *et al.* 1977, <sup>7</sup>Morton 1987, <sup>8</sup>Kirtikar & Basu 1988, <sup>9</sup>Anonymous 1952, <sup>10</sup>Harborne & Baxter 1999, \*Fruit and seed.

polyphenolic substances and flowers with numerous stamens (Landrum 1988). The family is divided into two sub-families, the Leptospermoideae and Myrtoideae. Edible fruits and useful spices, including guava, pitanga, clove and bay rum are produced by the closely related genera of the Myrtoideae, which can be divided into three subtribes. The subtribe Eugeniinae represents a large and morphologically diverse group, and most of these species have at one time been assigned to the genus *Eugenia*. Splitting the group depends on deciding which species to remove from *Eugenia* (McVaugh 1966), and the full taxonomic arrangement is still under debate (Mabberley 1993). They are pantropical in occurrence, concentrating in South America and Southeast Asia-Eastern Australia. The genus *Eugenia* is now mostly considered the neotropical group, numbering around 1000

species, and the plants designated *Syzygium* are generally considered by many to be the Old World genus. Other genera in this subtribe include *Myrciaria*, *Plinia*, *Catinga* and *Calycorectes*. Researchers disagree on the exact taxonomic arrangement, and many of these plants have multiple synonyms in at least one of these genera (Facciola 1998, McVaugh 1966, Popenoe 1920).

The fruits of this subtribe are often described by their bright anthocyanin colors, including orange, red, purple and black (dark purple). Examples of fruits in the subtribe are shown in Figure 2. They are sweet to tart, aromatic and many are astringent, indicating the presence of tannins. The taste is often described as somewhat acid. New shoot growth for many species is wine-colored (Facciola 1998, Popenoe



**Figure 2.** Examples of fruits in Myrtaceae subtribe Eugeniinae. **A.** *Eugenia brasiliensis*, **B.** *Eugenia uniflora*, **C.** *Myrciaria cauliflora*, **D.** *Syzygium cumini*.

1920), suggesting a high anthocyanin content. Many known antioxidant flavonoids have been isolated from all parts of many species in this group. Haron *et al.* found that flavonoid composition of New World and Old World *Eugenia* species is similar (Haron *et al.* 1992). Seventeen *Eugenia* species were tested, and myricetin was present in leaf extracts of 100%, quercetin in 71%, and kaempferol in 24%. Ellagic acid, procyanidin and prodelphinidin were found in most species tested, and Nair and Subramanian (Nair and Subramanian 1962) found that ellagic acid and myricetin are common throughout the family, as is methyl ellagic acid (Hegnauer 1990). In dicots, ellagic acid is usually confined to certain families and plants containing trihydroxy flavonoids, just as caffeic and p-coumeric acids are found along with corresponding di- and monohydroxy flavonoids (Bate-Smith 1963). Theoduloz showed that flavonoids in the five species of the Myrtaceae tested inhibit xanthine oxidase activity (Theoduloz *et al.* 1988), Schmeda-Hirschman credits this activity to the presence of the flavonoids quercitrin, quercetin, myricitrin, and myricetin (Schmeda-Hirschmann *et al.* 1987). High doses of the leaves showed no toxicity.

Surinam cherry (*E. uniflora*) is widely regarded as one of the best tasting of the *Eugenia* species, and the fruits average about 1 inch in size. They have a characteristic ribbed appearance, and several cultivars have been developed with fruits ranging from orange to crimson to black (Facciola 1998, Morton 1987). There is an extensive amount of literature documenting the ethnomedical uses of the leaves of Surinam cherry (Consolini *et al.* 1999, Duke 2000, Rücker *et al.* 1977, Schapoval *et al.* 1994, Schmeda-Hirschmann *et al.* 1987, Weyerstahl *et al.* 1988), and most of the phytochemical work has subsequently focused on characterizing the essential oil of the leaves (Weyerstahl *et al.* 1988). Ascorbic acid, b-carotene, and a few sesquiterpenes have been identified from the fruits (Duke 2000, Rücker *et al.* 1977), but no studies of flavonoid content of the fruit have been done. Several well-known antioxidant flavonoids have been reported from leaf extracts, including myricetin, myricitrin, gallic acid, quercetin, and quercitrin (Schmeda-Hirschmann *et al.* 1987) as well as the tannins eugeniflorin D-1 and D-2 (Lee *et al.* 1997). Popenoe notes that the Brazilians prepare a liqueur from the fruits, and consider syrups and wines to have a medicinal value (Popenoe 1920). In Ma-


deira, fruits of *E. uniflora* are eaten for intestinal troubles (Rivera & Obón 1995). Fruits and leaves are also used for their astringent qualities, and are active against high blood pressure (Bandoni *et al.* 1972). Water decoctions of *E. uniflora* leaves are used in Paraguay to lower cholesterol and blood pressure (Ferro *et al.* 1988), and have a highly significant anti-inflammatory action (Schapoval, *et al.* 1994). Ferro showed that leaf extracts were slightly active on lipid metabolism, and may exert a protective effect on triglycerides and very low-density lipoprotein levels (Ferro *et al.* 1988).

Other fruits in this subtribe are also colorful, with an extensive ethnobotanical and ethnomedical use that suggests a possible flavonoid content. *S. jambos* fruit is used as a tonic for the brain and for liver problems, as an astringent, and digestive and diuretic (Kirtikar & Basu 1988, Morton 1987). The leaves contain seventeen different flavonoids (Constant *et al.* 1997, Slowing, *et al.* 1996, Slowing *et al.* 1994) and are used as an anesthetic, anti-inflammatory, and astringent, for apoplexy, asthma, bronchitis, cough, diabetes, dysentery, influenza, and rheumatism (Rivera & Obón 1995). *S. samarangense*, which is cultivated in India for its edible fruit (Anonymous 1952) contains two flavonol glycosides as well as epigallocatechin 3-O-gallate, epicatechin 3-O-gallate, and samarangenin A and B (Harborne and Baxter 1999). In Taiwan, the flowers, which contain tannins, are used to treat fever and halt diarrhea. Flowers also contain desmethoxymatteucinol, 5-O-methyl-4'-desmethoxymatteucinol, oleanic acid and b-sitosterol (Morton 1987). The jaboticaba (*Myrciaria cauliflora*) is a popular edible in Brazil, much like grapes in the U.S. (Popenoe 1920). They are a dark red to maroon-purple and black, and are used to make jam, tarts, strong wine and a liqueur (Facciola 1998). *E. aggregata* is a popular reddish-purple edible in Brazil, eaten fresh or used to make jams and jellies (Facciola 1998) which has not been phytochemically examined. In preparing this study, the phytochemistry and ethnobotany of each plant was reviewed and noted, but those uses and compounds associated with the fruit only were given more emphasis, and are summarized in Table 1.

### Laboratory Work

In the current study, fruits were homogenized in a blender with methanol and extracted for one to two hours. Methanolic extracts were concentrated in vacuo and partitioned using a solvent-solvent procedure depicted in Figure 3. Hexane extracts contain lipid-soluble antioxidant vitamins such as b-carotene and a-tocopherol. Aqueous extracts contain sugars and ascorbic acid. The EtOAc fraction, being moderately polar, contains the polyphenolic components.


**Figure 3.** Partitioning scheme for Myrtaceous fruits.




EtOAc fractions were tested in a simple DPPH chemical assay following Yamaguchi (Yamaguchi *et al.* 1998). DPPH (Figure 4) produces a stable free radical, and is a good indicator of radical scavenging capacity. The 50% inhibition concentration (IC<sub>50</sub>) value is obtained using serial dilutions. A lower IC<sub>50</sub> value indicates greater activity. An IC<sub>50</sub> less than 50 µg/mL is considered very active, 50 – 100 is moderately active, 100 – 200 is slightly active and a value above 200 µg/mL is considered inactive. Ascorbic acid and a-tocopherol are used as positive controls, with IC<sub>50</sub> values of 18.3 µg/mL and 53.3 µg/mL, respectively.

The EtOAc fraction of *E. uniflora* was separated by vacuum-liquid chromatography and the most active fraction (SC<sub>10</sub>) was analyzed by high performance liquid chromatography (HPLC) using an on-line DPPH assay developed by Koleva (Koleva *et al.* 2000) and adopted in our laboratory (Figure 5). This assay demonstrates the antioxidant activity of individual phytochemicals. Spec-

**Figure 4.** Stable free radical, of DPPH that is a good indicator of radical scavenging capacity.





**Figure 5.** On-line Dpph-HPLC scheme.

tral data is collected both at 517 nm (for DPPH activity) and 230-500 nm (for flavonoid detection). After reacting, the radical absorbing activity is depicted on the chromatogram as a negative peak. As seen in Figure 6, the detection of compounds is graphically correlated to the activity in the assay. The main constituent eluted at 11.9 minutes and was very active in the assay as can be seen from the negative peak. The identity of this compound remains to be determined. Based on its UV absorbance profile it is very likely a flavonoid. The profile (Figure 6, Inset A) is very similar to quercitrin, which has been isolated from leaf extracts of *E. uniflora*.


## Conclusions and Discussion

Renowned pantropically as both food and medicine, these fruits have long been incorporated into traditional holistic health systems. Western medicine has, for the most part, overlooked them as potentially beneficial dietary components. Most of the pharmacological studies have not focused on the fruits. This study demonstrates that there are antioxidant compounds in these fruits. The most active fruits in the DPPH assay are *E. foetida* (15.9  $\mu\text{g}/\text{mL}$ ) and *E. uniflora* (19.6  $\mu\text{g}/\text{mL}$ ). *M. cauliflora* was also very active

at 35  $\mu\text{g}/\text{mL}$ . *E. aggregata*, *E. stipitata* and *S. samarangense* were moderately active, with IC<sub>50</sub> values of 74.1  $\mu\text{g}/\text{mL}$ , 79.0  $\mu\text{g}/\text{mL}$  and 76.8  $\mu\text{g}/\text{mL}$ , respectively. Only *S. jambos* appears to be inactive at 247  $\mu\text{g}/\text{mL}$ . These fruits compare favorably with known antioxidants ascorbic acid (18.3  $\mu\text{g}/\text{mL}$ ) and  $\alpha$ -tocopherol (53.3  $\mu\text{g}/\text{mL}$ ). Table 1 summarizes the DPPH assay results. Further understanding of the polyphenolic content of these fruits may be of great benefit in understanding the health aspects of both the traditional and modern uses of these fruits.

## Future Work

Active compounds from these species will be isolated and analyzed to determine chemical identity. Novel compounds will be elucidated using modern well-established analytical methods. Finally, the antioxidant capacity of beverages, vinegars and jams made from the most active fruits will be tested to see if the antioxidant compounds remained intact during the juicing and fermenting process.



**Figure 5.** Chromatogram of VLC fraction SC\_10 showing several active compounds in the on-line HPLC-DPPH assay. A negative peak at 517 nm corresponds to DPPH quenching, indicating an active compound. The UV absorbance spectrum of the major constituent is shown as inset A.

## Acknowledgements

Many thanks is given to those who have generously aided this project by allowing us to collect fruit: Larry Schokman, Director of The Kampong; Chris Rollins, Director of the Fruit and Spice Park; Dr. Jonathan H. Crane of the University of Florida Tropical Research and Education Center; and the Broward County Rare Fruit and Vegetable Council.

## Literature Cited

- Anonymous. 1952. *The Wealth of India*, Vol X, pp 93-107. Council of Scientific and Industrial Research, New Delhi.
- Bandoni, A.L., M.E. Mendiondo, R.V.D. Rondina & J.D. Coussio. 1972. Survey of Argentine medicinal plants. I. Folklore and phytochemical screening. *Lloydia* 35:69-80.
- Bate-Smith, E.C. 1963. Usefulness of chemistry in plant taxonomy as illustrated by the flavonoid constituents. Pp. 127-140 in *Chemical Plant Taxonomy*. Edited by T. Swain. Academic Press, London.
- Blumenthal, R.D., W. Lew, A. Reising, D. Soyne, L. Osorio, Z. Ying & D. Goldenberg. 2000. Antioxidant vitamins reduce normal tissue toxicity induced by radio-immunotherapy. *International Journal of Cancer* 86:276-280.
- Burns, J., P. Gardner, J. O'Neil, S. Crawford, I. Morecroft, D.B. McPhail, C. Lister, D. Matthews, M.R. MacLean, M.E.J. Lean, G.G. Duthie & A. Crozier. 2000. Relationship among antioxidant activity, vasodilation capacity, and phenolic content of red wines. *Journal of Agricultural Food Chemistry* 48:220-230.
- Cao, G., S.L. Booth, J.A. Sadowski & R. Prior. 1998. Increases in human plasma antioxidant capacity after consumption of controlled diets high in fruits and vegetables. *American Journal of Clinical Nutrition* 68:1081-1087.
- Cao, G., E. Sofic & R. Prior. 1997. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Radical Biology and Medicine* 22:749-760.
- Consolini, A.E., O.A. Baldini & A.G. Amat. 1999. Pharmacological basis for the empirical use of *Eugenia uniflora* L. (Myrtaceae) as antihypertensive. *Journal of Ethnopharmacology* 66:33-39.
- Constant, H.L., K. Slowing, J.G. Graham, J.M. Pezzuto, G.A. Cordell & C.W. Beecher. 1997. A general method for the dereplication of flavonoid glycosides utilizing high performance liquid chromatography/mass spectrometric analysis. *Phytochemical Analysis* 8:176-180.
- Dewick, P. 1997. *Medicinal Natural Products: A Biosynthetic Approach*. John Wiley & Sons, New York.
- Duke, J. 2000. Chemicals in *Eugenia uniflora*. In *Dr. Duke's Phytochemical and Ethnobotanical Databases: Agricultural Research Service, USDA*, <http://www.ars-grin.gov/duke>.
- Facciola, S. 1998. *Cornucopia II*. Kampong Publications, Vista, California.
- Ferro, E., A. Schinini, M. Maldonado, J. Rosner & G.S. Hirschman. 1988. *Eugenia uniflora* leaf extract and lipid me-

- tabolism in *Cebus apella* monkeys. *Journal of Ethnopharmacology* 24:321-325.
- Formica, J.V. & W. Regelson. 1995. Review of the biology of quercetin and related bioflavonoids. *Food and Chemical Toxicology* 33:1061-1080.
- Franco, M.R.B. & T. Shibamoto. 2000. Volatile composition of some Brazilian Fruits: Umbu-caja (*Spondias citherea*), camu-camu (*Myrciaria dubia*), araca-boi (*Theobroma grandiflorum*). *Journal of Agricultural Food Chemistry* 48:1263-1265.
- Frankel, E.N., J. Kanner, J.B. German, E. Parks & J.E. Kinsella. 1993a. Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *The Lancet* 341:454-457.
- Frankel, E.N., A.L. Waterhouse & J.E. Kinsella. 1993b. Inhibition of human LDL oxidation by resveratrol. *The Lancet* 341:1103-1104.
- Gibbs, R.D. 1993. History of Chemical Taxonomy. Pp. 41-88 in *Chemical Plant Taxonomy*. Edited by T. Swain. Academic Press, London.
- Giugliano, D. 2000. Dietary antioxidants for cardiovascular prevention. *Nutrition, Metabolism and Cardiovascular Disease* 10:38-44.
- Haeghele, A.D., C. Gillette, C. O'Neill, P. Wolfe, J. Heimedinger, S. Sedlacek & H.J. Thompson. 2000. Plasma xanthophyll carotenoids correlate inversely with indices of oxidative DNA damage and lipid peroxidation. *Cancer Epidemiology, Biomarkers and Prevention* 9:421-425.
- Harborne, J.B. 1963. Distribution of anthocyanins in higher plants. Pp. 359-388 in *Chemical Plant Taxonomy*. Edited by T. Swain. Academic Press, London.
- Harborne, J.B. & H. Baxter. 1999. Pages ix-xv, 30, 36, 39, 381, 384, 476 in *The Handbook of Natural Flavonoids*, Vol 2. John Wiley and Sons, New York.
- Haron, N.W., D.M. Moore & J.B. Harborne. 1992. Distribution and taxonomic significance of flavonoids in the genus *Eugenia* (Myrtaceae). *Biochemical Systematics and Ecology* 20:226-268.
- Hegnauer, R. 1990. *Chemotaxonomie der Pflanzen*, Vol. 9:119-129. Birkhäuser Verlag, Berlin.
- Hertog, M.G.L., E.J.M. Feskens, P. Hollman, M.B. Katan & D. Kromhout. 1993. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen elderly study. *The Lancet* 342:1007-1011.
- Hertog, M.G.L., D. Kromhout, C. Aravanis, H. Blackburn, R. Buzina, F. Fidanza, S. Glampaoli, A. Jansen, A. Menotti, S. Nedeljkovic, M. Pekkainen, B.S. Simic, H. Toshima, E.J.M. Feskens, P.C.H. Hollman & M.B. Katan. 1995. Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Archives of Internal Medicine* 155:381-386.
- Hollman, P.C. & M.B. Katan. 1999. Dietary flavonoids: intake, health effects and bioavailability. *Food and Chemical Toxicology* 37:937-942.
- Jacob, R.A. & B.J. Burri. 1996. Oxidative damage and defense. *American Journal of Clinical Nutrition* 63:985S-990S.
- Kähkönen, M., A. Hopia, H. Vuorela, J. Rauha, K. Pihlaja, T. Kujala & M. Heinonen. 1999. Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural Food Chemistry* 47:3954-3962.
- Kirtikar, K.R. & B.D. Basu. 1988. *Indian Medicinal Plants*, Vol. 2:1038-1063. International Book Distributors, Dehradun.
- Knekt, P., R. Järvinen, A. Reunanen & J. Maatela. 1996. Flavonoid intake and coronary mortality in Finland: a cohort study. *British Journal of Medicine* 312:478-481.
- Koleva, I.I., H.A.G. Niederlander & T.A. van Beck. 2000. An on-line HPLC method for detection of radical scavenging compounds in complex mixtures. *Analytical Chemistry* 72:2323-2328.
- Landrum, L.R. 1988. The Myrtle family (Myrtaceae) in Chile. *Proceedings of the California Academy of Science*. 1988:277-317.
- Lee, M.-H., S. Nishimoto, L.-L. Yang, K.-Y. Yen, T. Hatano, T. Yoshida & T. Okuda. 1997. Two macrocyclic hydrolyzable tannin dimers from *Eugenia uniflora*. *Phytochemistry* 44:1343-1349.
- Mabberley, D.J. 1993. *The Plant Book*. Bath Press, Avon.
- McVaugh, R. 1966. Tropical American Myrtaceae: notes on generic concepts and descriptions of previously unrecognized species. *Fieldiana: Botany* 29:145-228.
- Middleton, E. & S. Kandaswami. 1993. The impact of plant flavonoids on mammalian biology: implications for immunity, inflammation and cancer. Pp. 619-652 in *The Flavonoids: Advances in Research Since 1986*. Edited by J.B. Harborne. Chapman and Hall, London.
- Morton, J. 1987. *Fruits of Warm Climates*. Julia Morton, Winterville, North Carolina.

- Nair, A.G.R. & S.S. Subramanian. 1962. Chemical examination of the flowers of *Eugenia jambolana*. *Journal of Science and Industrial Research* 21B:457-458.
- Ness, A.R. & J.W. Powles. 1999. The role of diet, fruit and vegetables and antioxidants in the aetiology of stroke. *Journal of Cardiovascular Risk* 6:229-234.
- Pellegrini, N., P. Riso & M. Porrinni. 2000. Tomato consumption does not affect the total antioxidant capacity of plasma. *Nutrition* 16:268-271.
- Pierpoint, W.S. 1986. Flavonoids in the human diet. Pp. 125-140 in *Plant Flavonoids in Biology and Medicine: Biochemistry, Pharmacology, and Structure-Activity Relationship*. Edited by V. Cody, E. Middleton & J.B. Harborne. Alan R Liss, New York.
- Pietta, P.-G. 2000. Flavonoids as antioxidants. *Journal of Natural Products* 63:1035-1042.
- Popenoe, W. 1920. *Manual of Tropical and Subtropical Fruits*. Hafner Press, New York.
- Rice-Evans, C., N. Miller & G. Paganga. 1996. Structure-antioxidant activity relationships of flavonoids and phenolic compounds. *Free Radical Biology and Medicine* 20:933-956.
- Rice-Evans, C.A., N.J. Miller & G. Paganga. 1997. Antioxidant properties of phenolic compounds. *Trends in Plant Science* 2:152-159.
- Rivera, D. & C. Obón. 1995. The ethnopharmacology of Madeira and Porto Santo islands, a review. *Journal of Ethnopharmacology* 46:73-93.
- Robinson, T. 1991. *The Organic Constituents of Higher Plants*. Cordu Press, North Amherst.
- Rücker, G., G.A.A.B. Silva, L. Bauer & M. Schikarski. 1977. New constituents of *Stenocalyx michelii*. *Planta Medica* 31:322.
- Schapoval, E.E., S.M. Silveira, M.L. Miranda, C.B. Alice & A.T. Henriques. 1994. Evaluation of some pharmacological activities of *Eugenia uniflora* L. *Journal of Ethnopharmacology* 44:137-142.
- Scheen, A.J. 2000. Antioxidant vitamins in the prevention of cardiovascular disease. Second part: results of clinical trials. *Revue Medicale de Liege* 55:105-109.
- Schmeda-Hirschmann, G., C. Theoduloz, L. Franco, E. Ferro & A. Rojas De Arias. 1987. Preliminary pharmacological studies on *Eugenia uniflora* leaves: xanthine oxidase inhibitory activity. *Journal of Ethnopharmacology* 21:183-186.
- Silva, C.T.C. 1997. Postharvest modifications in camucamu fruit (*Myrciaria dubia* McVaugh) in response to stage of maturation and modified atmosphere. Pp. 23-26 in *International Symposium on Myrtaceae; International Society for Horticultural Sciences*. Curitiba, Brazil.
- Slowing, K., E. Carretero & A. Villar. 1996. Anti-inflammatory compounds of *Eugenia jambos*. *Phytotherapy Research* 10:S126-S127.
- Slowing, K., M. Sollhuber, E. Carretero & A. Villar. 1994. Flavonoid glycosides from *Eugenia jambos*. *Phytochemistry* 37:255-258.
- Steinberg, D. 1997. A critical look at the evidence for the oxidation of LDL in atherogenesis. *Atherosclerosis* 131(Suppl.):S5-S7.
- Theoduloz, C., L. Franco, E. Ferro & G. Schmeda-Hirschman. 1988. Xanthine oxidase inhibitory activity of Paraguayan Myrtaceae. *Journal of Ethnopharmacology* 24:179-183.
- van Poppel, G., A. Kardinaal, H. Princen & F.J. Kok. 1994. Antioxidants and coronary heart disease. *Annals of Medicine* 26:429-434.
- Vinson, J., J. Jang, J. Yang, Y. Dabbagh, X. Liang, M. Serry, J. Proch & S. Cai. 1999a. Vitamins and especially flavonoids in common beverages are powerful in vitro antioxidants which enrich low density lipoproteins and increase their oxidative resistance after ex vivo spiking in human plasma. *Journal of Agricultural Food Chemistry* 47:2502-2504.
- Vinson, J., J. Proch & L. Zubik. 1999b. Phenol antioxidant quantity and quality in foods: cocoa, dark chocolate, and milk chocolate. *Journal of Agricultural Food Chemistry* 47:4821-4824.
- Wang, H., G. Cao & R. Prior. 1997. Oxygen radical absorbing capacity of anthocyanins. *Journal of Agricultural Food Chemistry* 45:304-309.
- Weyerstahl, P., H. Marschall-Weyerstahl, C. Christiansen, B.O. Oguntimein & A.O. Adeoye. 1988. Volatile constituents of *Eugenia uniflora* leaf oil. *Planta Medica* 54:546-549.
- Yamaguchi, T., H. Takamura, T. Matoba & J. Terao. 1998. HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl. *Bioscience, Biotechnology and Biochemistry* 62:1201-1204.
- Yoshikawa, M., H. Shimada, N. Nishida, Y. Li, I. Toguchida, J. Yamahara & H. Matsuda. 1998. Antidiabetic principles of natural medicines. II. Aldose reductase and a-

glucosidase inhibitors from Brazilian natural medicine, the leaves of *Myrcia multiflora* DC. (Myrtaceae): Structures of myrciacitrins I and II and myrciaphenones A and B. *Chemical and Pharmaceutical Bulletin* 46:113-119.

