

# Index to Volume 28

## Author Index

- ALLEN, GERALD R.:  
A review of the labrid genus *Paracheilinus*, with the description of a new species from Melanesia, 449-455
- ARVIK, JON H., and DAN L. WILLSON:  
Soil algae of Eniwetok Atoll, the Marshall Islands, 189-190
- BANNER, ALBERT H., and DORA M. BANNER:  
Contributions to the knowledge of the alpheid shrimp of the Pacific Ocean. Part XVII. Additional notes on the Hawaiian alpheids: new species, subspecies, and some nomenclatorial changes, 423-437
- BANNER, DORA M.:  
*see* Banner and Banner
- BRADFORD, JANET M.:  
*Euchaeta marina* (Prestandrea) (Copepoda, Calanoida) and two closely related new species from the Pacific, 159-169
- BURGESS, WARREN E.:  
Evidence for the elevation to family status of the angelfishes (Pomacanthidae), previously considered to be a subfamily of the butterflyfish family, Chaetodontidae, 57-71
- CASSIE, VIVIENNE:  
Algal flora of some North Island, New Zealand lakes, including Rotorua and Rotoiti, 467-505
- CHAVE, E. H., and D. B. ECKERT:  
Ecological aspects of the distributions of fishes at Fanning Island, 297-317
- COHEN, DANIEL M.:  
The ophidioid fish genus *Luciobrotula* in the Hawaiian Islands, 109-110
- COLES, STEPHEN L.:  
*see* Jokiel and Coles
- COLLETTE, BRUCE B.:  
Geographic variation in the central Pacific halfbeak *Hyporhamphus acutus* (Günther), 111-122
- DUNN, DAPHNE FAUTIN:  
*Actinogeton sesere* (Coelenterata, Actiniaria) in Hawaii, 181-188  
*Radianthus papillosa* (Coelenterata, Actiniaria) re-described from Hawaii, 171-179  
Redescription of *Anthopleura nigrescens* (Coelenterata, Actiniaria) from Hawaii, 377-382
- EDGERLEY, W. H. L.:  
*see* Glasby and Edgerley
- EGLOFF, J.:  
*see* Johnson, Egloff, and Hemler
- FELL, F. JULIAN:  
The echinoids of Easter Island (Rapa Nui), 147-158
- FIELD, LAURENCE H.:  
A description and experimental analysis of Batesian mimicry between a marine gastropod and an amphipod, 439-447
- FOSTER, B. A.:  
The barnacles of Fiji, with observations on the ecology of barnacles on tropical shores, 35-56
- GLASBY, G. P., and W. H. L. EDGERLEY:  
Geochemistry of lake waters from the South Island, New Zealand, 507-515
- GLYNN, CARMEN S.:  
*see* Glyn and Glynn
- GLYNN, PETER W., and CARMEN S. GLYNN:  
On the systematics of *Ancinus* (Isopoda, Sphaeromatidae), with the description of a new species from the tropical eastern Pacific, 401-422
- GUINThER, ERIC B.:  
Groundwater and nearshore hyposaline conditions at Fanning Island during a period of higher than normal rainfall, 319-337
- HEMLER, L. G.:  
*see* Johnson, Egloff, and Hemler
- HUMES, ARTHUR G.:  
Cyclopoid copepods associated with the coral genera *Favia*, *Favites*, *Platygyra*, and *Merulina* in New Caledonia, 383-399
- JOHNSON, G. L., J. EGLOFF, and L. G. HEMLER:  
Geophysical observations between Hawaii and Australia, 191-205
- JOKIEL, PAUL L., and STEPHEN L. COLES:  
Effects of heated effluent on hermatypic corals at Kahe Point, Oahu, 1-18  
*see also* Jokiel and Townsley
- JOKIEL, PAUL L., and SIDNEY J. TOWNSLEY:  
Biology of the polyclad *Prosthiostomum* (*Prosthiostomum*) sp., a new coral parasite from Hawaii, 361-373  
*see also* Jokiel and Coles
- KAY, E. ALISON:  
Fanning Island: editor's note, 207-210  
*see also* Kay and Switzer
- KAY, E. ALISON and MARILYN F. SWITZER:  
Molluscan distribution patterns in Fanning Island Lagoon and a comparison of the mollusks of the lagoon and the seaward reefs, 275-295  
*see also* Kay, E. Alison
- KAY, JOHN C.:  
*see* RANDALL and KAY

- MARAGOS, J. E.:  
Coral communities on a seaward reef slope, Fanning Island, 257-273  
Reef corals of Fanning Island, 247-255
- MARGOLIS, S. V.:  
*see* Williams and Margolis
- V. MEJSTRÍK:  
The frequency of vesicular-arbuscular mycorrhizae in the roots of *Camellia japonica* L. from different sites in New Zealand, 73-77
- MEYERS, GARY A.:  
*see* Stroup and Meyers
- MORTON, BRIAN:  
Some aspects of the biology, population dynamics, and functional morphology of *Musculista senhousia* Benson (Bivalvia, Mytilidae), 19-33
- PESRET, F.:  
*see* Smith and Pesret
- RANDALL, JOHN E., and JOHN C. KAY:  
*Stethojulis axillaris*, a junior synonym of the Hawaiian labrid fish *Stethojulis balteata*, with a key to the species of the genus, 101-107
- ST. JOHN, HAROLD:  
A new *Ervatamia* (Apocynaceae) from Makatea Island, Tuamotu Archipelago. Pacific Plant Studies 23, 436-465  
Revision of the genus *Pandanus* Stickman. Part 36, The new section *Asperi* from Indomalaya, 79-82
- Revision of the genus *Pandanus* Stickman. Part 37, *Pandanus* on Aldabra Island, Indian Ocean, 83-100  
*Skottsbergiliana* new genus (Cucurbitaceae) of Hawaii Island. Hawaiian plant studies 41, 457-462.  
The vascular flora of Fanning Island, Line Islands, Pacific Ocean, 339-355
- SASSAMAN, CLAY:  
Capacity for development of secondary manubria in *Eutonina indicans* medusae (Hydrozoa), 375-376
- SMITH, S. V., and F. PESRET:  
Processes of carbon dioxide flux in the Fanning Island Lagoon, 225-245
- STROUP, EDWARD D., and GARY A. MEYERS:  
The flood-tide jet in Fanning Island Lagoon, 211-223
- TOWNSLEY, SIDNEY J.:  
*see* Jokiel and Townsley
- WILLIAMS, J. A., and S. V. MARGOLIS:  
Sipunculid burrows in coral reefs: evidence for chemical and mechanical excavation, 357-359
- WILLSON, DAN L.:  
*see* Arvik and Willson
- YAMAGUCHI, MASASHI:  
Effect of elevated temperature on the metabolic activity of the coral reef asteroid *Acanthaster planci* (L.), 139-146  
Growth of juvenile *Acanthaster planci* (L.) in the laboratory, 123-138

### Subject Index\*

- Acanthaster planci* (L.)  
effect of elevated temperature on the metabolic activity of, 139-146  
growth of juvenile, in the laboratory, 123-138
- Aldabra Island  
revision of the genus *Pandanus* Stickman. Part 37, *Pandanus* on, Indian Ocean, 83-100
- Algae  
of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467-505  
soil, of Eniwetok Atoll, the Marshall Islands, 189-190
- Alpheus heeia* sp. nov., 423-437  
*Alpheus lobidens polynesica* subsp. nov., 423-437  
*Alpheus mackayi* nov. comb., 423-437
- Amphineura  
molluscan distribution patterns in Fanning Island Lagoon and a comparison of the mollusks of the lagoon and the seaward reefs, 275-295
- Ancinus panamensis* n. sp., 401-422
- Angelfishes, evidence for the elevation to family status of the, 57-71
- Angiospermae  
frequency of vesicular-arbuscular mycorrhizae in the roots of *Camellia japonica* L. from different sites in New Zealand, 73-77  
a new *Ervatamia* (Apocynaceae) from Makatea Island, Tuamotu Archipelago. Pacific plant studies 23, 463-465  
revision of the genus *Pandanus* Stickman. Part 36, the new section *Asperi* from Indomalaya, 79-82  
revision of the genus *Pandanus* Stickman. Part 37, *Pandanus* on Aldabra Island, Indian Ocean, 83-100  
*Skottsbergiliana* new genus (Cucurbitaceae) of Hawaii Island. Hawaiian plant studies 41, 457-462  
vascular flora of Fanning Island, Line Islands, Pacific Island, 339-355
- Anthozoa  
*Actinogeton sesere* in Hawaii, 181-188  
biology of the polyclad *Prosthiostomum* (*Prosthiostomum*) sp., a new coral parasite from Hawaii, 361-373  
coral communities on a seaward reef slope, Fanning Island, 257-273

\* Biological organisms are indexed under their respective classes.

- Anthozoa (*cont.*)  
 cyclopoid copepods associated with the coral genera *Favia*, *Favites*, *Platygyra*, and *Merulina* in New Caledonia, 383–399  
 effects of heated effluent on hermatypic corals at Kahe Point, Oahu, 1–18  
*Radianthus papillosa* redescribed from Hawaii, 171–179  
 redescription of *Anthopleura nigrescens* from Hawaii, 377–382  
 reef corals of Fanning Island, 247–255  
 sipunculid burrows in coral reefs: evidence for chemical and mechanical excavation, 357–359
- Asperi* sect. nov., 79–82
- Astrozoidea  
 effect of elevated temperature on the metabolic activity of the coral reef asteroid *Acanthaster planci* (L.), 139–136  
 growth of juvenile *Acanthaster planci* (L.) in the laboratory, 123–138
- Australia  
 geophysical observations between Hawaii and Australia, 191–205
- Bacillariophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505
- Barnacles, ecology of, on tropical shores, 35–56
- Batesian mimicry, a description and experimental analysis of, between a marine gastropod and an amphipod, 439–447
- Bivalvia  
 molluscan distribution patterns in Fanning Island Lagoon and a comparison of the mollusks of the lagoon and the seaward reefs, 275–295  
 some aspects of the biology, population, dynamics, and functional morphology of *Musculista senhousia* Benson (Bivalvia, Mytilidae), 19–33
- Camellia japonica*, frequency of vesicular-arbuscular mycorrhizae in roots of, 73–77
- Chlorophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505
- Chrysophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505  
 soil algae of Eniwetok Atoll, the Marshall Islands, 189–190
- Contributions to the knowledge of the alpheid shrimp of the Pacific Ocean  
 XVII. Additional notes on the Hawaiian alpheids: new species, subspecies, and some nomenclatorial changes, 423–437
- Coral reefs, sipunculid burrows in, 357–359
- Corals  
 hermatypic, effects of heated effluent on, 1–18  
 reef, of Fanning Island, 247–255
- Crustacea  
 barnacles of Fiji, with observations on the ecology of barnacles on tropical shores, 35–56
- Contributions to the knowledge of the alpheid shrimp of the Pacific Ocean. Part XVII. Additional notes on the Hawaiian alpheids: new species, subspecies, and some nomenclatorial changes, 423–437
- cyclopoid copepods associated with the coral genera *Favia*, *Favites*, *Platygyra*, and *Merulina* in New Caledonia, 383–399  
 a description and experimental analysis of Batesian mimicry between a marine gastropod and an amphipod, 439–447  
*Eucbaeta marina* (Prestandrea) (Copepoda, Calanoida) and two closely related new species from the Pacific, 159–169  
 on the systematics of *Ancinus* (Isopoda, Sphaeromatidae), with the description of a new species from the tropical eastern Pacific, 401–422
- Cryptophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505
- Dinophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505
- Easter Island  
 echinoids of, 147–158
- Echinozoa  
 echinoids of Easter Island (Rapa Nui), 147–158
- Eniwetok Atoll  
 soil algae of, 189–190  
*Ervatamia makateensis* sp. nov., 463–465  
*Eucbaeta marinella* sp. n., 159–169  
*Eucbaeta rimana* sp. n., 159–169
- Euglenophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505
- Eutonina indicans*, capacity for development of secondary manubria in medusae of, 375–376
- Fanning Island  
 coral communities on a seaward reef slope at, 257–273  
 ecological aspects of the distributions of fishes at, 297–317  
 editor's note, 207–210  
 flood-tide jet in lagoon at, 211–223  
 groundwater and nearshore hyposaline conditions at, during a period of higher than normal rainfall, 319–337  
 molluscan distribution patterns in lagoon and a comparison of the mollusks of the lagoon and the seaward reefs at, 275–295  
 processes of carbon dioxide flux in the lagoon at, 225–245  
 reef corals of, 247–255  
 vascular flora of, 339–355
- Fiji  
 barnacles of, with observations on the ecology of barnacles on tropical shores, 35–56

## Gastropoda

- a description and experimental analysis of Batesian mimicry between a marine gastropod and an amphipod, 439-447
- molluscan distribution patterns in Fanning Island Lagoon and a comparison of the mollusks of the lagoon and the seaward reefs, 275-295

## Hawaiian Islands

- Actiniogeton sesere* (Coelenterata, Actiniaria) in, 181-188
- biology of the polyclad *Prosthiostomum* (*Prosthiostomum*) sp., a new coral parasite from, 361-373
- effects of heated effluent on hermatypic corals at Kahe Point, Oahu, 1-18
- geophysical observations between Hawaii and Australia, 191-205
- the ophiroid fish genus *Luciobrotula* in, 109-110
- Radianthus papillosa* (Coelenterata, Actiniaria) re-described from, 171-179
- redescription of *Anthopleura nigrescens* (Coelenterata, Actiniaria) from, 377-382
- Skottsbergiliana* new genus (Cucurbitaceae) of Hawaii Island. Hawaiian plant studies 41, 457-462
- Stethojulis axillaris*, a junior synonym of the Hawaiian labrid fish *Stethojulis balteata*, with a key to the species of the genus, 101-107

## Hawaiian plant studies

- 41. *Skottsbergiliana* new genus (Cucurbitaceae) of Hawaii Island, 457-462

## Hong Kong

- some aspects of the biology, population dynamics, and functional morphology of *Musculista senhousia* Benson (Bivalvia, Mytilidae), 19-33

## Hydrozoa

- capacity for development of secondary manubria in *Eutonia indicans* medusae, 375-376
- Hyporhamphus acutus* (Günther), geographic variation in, in the central Pacific, 111-122

## Indomalaya

- revision of the genus *Pandanus* Stickman. Part 36, the new section of *Asperi* from, 79-82

## Key

- to the genera of Fijian balanomorphs, 39
- to the genus *Luciobrotula*, 109
- to sections and species of *Pandanus* on Aldabra Island, Indian Ocean, 83
- to the species of *Ancinus*, 417-418
- to the species of *Paracheilinus*, 450
- to the species of *Skottsbergiliana*, 457
- to the species of *Stethojulis*, 102-103

## Lamellibranchia. See Bivalvia

- Leptalphus pacificus* sp. nov., 423-437
- Luciobrotula* in the Hawaiian Islands, 109-110

## Marshall Islands

- soil algae of Eniwetok Atoll, 189-190
- Musculista senhousia* Benson, some aspects of the biol-

ogy, population dynamics, and functional morphology of, 19-33

- Mycorrhizae, vesicular-arbuscular, in roots of *Camellia japonica*, 73-77
- Myxophyceae
  - algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467-505
  - soil algae of Eniwetok Atoll, the Marshall Islands, 189-190

## New Caledonia

- cyclopoid copepods associated with the coral genera, *Favia*, *Favites*, *Platygyra*, and *Merulina* in, 383-399
- New combination
  - Alpheus mackayi* (Crustacea), 423-437
  - Skottsbergiliana lasiocephala* (Angiospermae), 457-462
- New cultivar
  - Pandanus tectorius* Warb., var. *novi-caledonicus* Martelli, cv. 'Kaina' (Angiospermae), 339-355

## New genus

- Skottsbergiliana* (Angiospermae), 457-462

## New Guinea

- a review of the labrid genus *Paracheilinus*, with the description of a new species from Melanesia, 449-455

## New section

- Asperi* (Angiospermae), 79-82

## New species

- Ancinus panamensis* (Crustacea), 401-422
- Alpheus heeia* (Crustacea), 423-437
- Ervatamia makateaensis* (Angiospermae), 463-465
- Eucbaeta marinella* (Crustacea), 159-169
- Eucbaeta rimana* (Crustacea), 159-169
- Leptalphus pacificus* (Crustacea), 423-437
- Pandanus aldabraensis* (Angiospermae), 83-100
- Pandanus alloios* (Angiospermae), 83-100
- Pandanus asper* (Angiospermae), 79-82
- Pandanus cbelyon* (Angiospermae), 83-100
- Pandanus fanningensis* (Angiospermae), 339-355
- Pandanus impar* (Angiospermae), 83-100
- Pandanus intraconicus* (Angiospermae), 83-100
- Pandanus subcubicus* (Angiospermae), 83-100
- Paracheilinus filamentosus* (Pisces), 449-455
- Skottsbergiliana partita* (Angiospermae), 457-462

## New subspecies

- Alpheus lobidens polynesica* (Crustacea), 423-437

## New Zealand

- algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467-505
- frequency of vesicular-arbuscular mycorrhizae in the roots of *Camellia japonica* L. from different sites in, 73-77
- geochemistry of lake waters from the South Island, 507-515

## Pacific plant studies

- 23. a new *Ervatamia* (Apocynaceae) from Makatea Island, Tuamotu Archipelago, 463-465
- Pandanus aldabraensis* sp. nov., 83-100
- Pandanus alloios* sp. nov., 83-100
- Pandanus asper* sp. nov., 79-82
- Pandanus cbelyon* sp. nov., 83-100

- Pandanus fanningensis* sp. nov., 339–355  
*Pandanus impar* sp. nov., 83–100  
*Pandanus intraconicus* sp. nov., 83–100  
*Pandanus subcubicus* sp. nov., 83–100  
*Pandanus tectorius* Warb., var. *novi-caledonicus* Martelli, cv. 'Kaina,' cultivar nov., 339–355  
*Paracheilinus filamentosus* n. sp., 449–455
- Phycomycetes  
 frequency of vesicular-arbuscular mycorrhizae in the roots of *Camellia japonica* L. from different sites in New Zealand, 73–77
- Pisces  
 geographic variation in the central Pacific halfbeak *Hyporhamphus acutus* (Günther), 111–122  
 ophidioid fish genus *Luciobrotula* in the Hawaiian Islands, 109–110  
 a review of the labrid genus *Paracheilinus*, with the description of a new species from Melanesia, 449–455  
 ecological aspects of the distributions of fishes at Fanning Island, 297–317  
 evidence for the elevation to family status of the angel-fishes (Pomacanthidae), previously considered to be a subfamily of the butterflyfish family, Chaetodontidae, 57–71  
*Stethojulis axillaris*, a junior synonym of the Hawaiian labrid fish *Stethojulis balteata*, with a key to the species of the genus, 101–107
- Pomacanthidae, evidence for the elevation to family status of the, 57–71
- Revision of the genus *Pandanus* Stickman  
 36. the new section *Asperi* from Indomalaya, 79–82  
 37. *Pandanus* on Aldabra Island, Indian Ocean, 83–100
- Sipuncula  
 sipunculid burrows in coral reefs: evidence for chemical and mechanical excavation, 357–359  
*Skottsbergiliana* gen. nov., 457–462  
*Skottsbergiliana lasiocephala* comb. nov., 457–462  
*Skottsbergiliana partita* sp. nov., 457–462
- Solomon Islands  
 a review of the labrid genus *Paracheilinus*, with the description of a new species from Melanesia, 449–455
- Tuamotu Archipelago  
 a new *Ervatamia* (Apocynaceae) from Makatea Island, 463–465
- Turbellaria  
 biology of the polyclad *Prosthiostomum* (*Prosthiostomum*) sp., a new coral parasite from Hawaii, 361–373
- Xanthophyceae  
 algal flora of some North Island, New Zealand, lakes, including Rotorua and Rotoiti, 467–505